


PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>7</sup> : <b>C07D 241/26, A61K 31/4965</b>		A1	(11) International Publication Number: <b>WO 00/12488</b>
			(43) International Publication Date: 9 March 2000 (09.03.00)
(21) International Application Number: PCT/EP99/06248 (22) International Filing Date: 26 August 1999 (26.08.99) (30) Priority Data: 9818881.6                      28 August 1998 (28.08.98)                      GB (71) Applicant (for all designated States except US): GLAXO GROUP LIMITED [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB). (72) Inventors; and (75) Inventors/Applicants (for US only): COX, Brian [GB/GB]; Glaxo Wellcome plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). HEALY, Mark, Patrick [GB/GB]; Glaxo Wellcome plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). WILD, Deborah [GB/GB]; Glaxo Wellcome plc, Gunnels Wood Road, Stevenage, Hertfordshire SG1 2NY (GB). (74) Agent: VOLCKMAN, Janis, F.; Glaxo Wellcome plc, Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(54) Title: PYRAZINE COMPOUNDS  <div style="text-align: center;">  <p>(I)</p> </div>			
(57) Abstract  A compound of formula (I) wherein R <sup>1</sup> is phenyl substituted by one or more halogen atoms; R <sup>2</sup> is -NH <sub>2</sub> ; R <sup>3</sup> is -NH <sub>2</sub> or hydrogen; R <sup>4</sup> is -CXNR <sup>a</sup> R <sup>b</sup> , -CXNH-(CH <sub>2</sub> ) <sub>y</sub> -NR <sup>a</sup> R <sup>b</sup> , wherein X is = O or = S; y is an integer zero, 1 or 2; R <sup>a</sup> and R <sup>b</sup> , which may be the same or different, are selected from hydrogen and C <sub>1-4</sub> alkyl or together with the nitrogen atom to which they are attached form a saturated 5- or 6-membered heterocycle containing one or two nitrogen heteroatoms, which heterocycle can be further substituted with one or more C <sub>1-4</sub> alkyl groups; and pharmaceutically acceptable derivatives thereof.			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		


## PYRAZINE COMPOUNDS

The present invention relates to a class of pyrazine compounds which are useful in the treatment of central nervous system (CNS) diseases and disorders and to their pharmaceutically acceptable derivatives, to pharmaceutical compositions containing them, to their use in the treatment of such disorders and to methods of preparation.

Numerous phenyl pyrazine derivatives are known in the prior art. For example, Synthesis (1987), (10), 914-915, describes phenyl pyrazine derivatives including, inter alia, 3-(4-chlorophenyl)-pyrazinamine. No pharmaceutical utility is however described in that prior art document.

The present invention relates to pyrazine derivatives which are sodium channel blockers. The compounds are surprisingly potent anti-convulsants having increased potency with respect to lamotrigine and increased selectivity in terms of CNS side-effects and inhibition of the enzyme dihydrofolate reductase. The compounds are therefore useful in the treatment of CNS diseases such as epilepsy.

Accordingly, the present invention provides a compound of formula (I)



wherein

R<sup>1</sup> is phenyl substituted by one or more halogen atoms;

$R^2$  is  $-NH_2$ ;

$R^3$  is  $-NH_2$  or hydrogen;

5  $R^4$  is  $-CXNR^aR^b$ ,  $-CXNH-(CH_2)_y-NR^aR^b$ ;

wherein

X is  $=O$  or  $=S$ ;

10

y is an integer zero, 1 or 2;

15  $R^a$  and  $R^b$ , which may be the same or different, are selected from hydrogen and  $C_{1-4}$  alkyl, or together with the nitrogen atom to which they are attached, form a saturated 5- or 6 - membered heterocycle containing one or two nitrogen heteroatoms, which heterocycle can be further substituted with one or more  $C_{1-4}$  alkyl groups;

and pharmaceutically acceptable derivatives thereof.

20

By pharmaceutically acceptable derivative is meant any pharmaceutically acceptable salt, solvate or ester, or salt or solvate of such ester of the compounds of formula (I), or any other compound which upon administration to the recipient is capable of providing (directly or indirectly) a compound of formula (I) or an active metabolite or residue thereof (eg. a prodrug).

25

It will be appreciated that, for pharmaceutical use, the salts referred to above will be the physiologically acceptable salts, but other salts may find use, for example in the preparation of compounds of formula (I) and the physiologically acceptable salts thereof.

30

Suitable pharmaceutically acceptable salts of the compounds of formula (I) include acid addition salts formed with inorganic or organic acids, preferably inorganic acids, e.g. hydrochlorides, hydrobromides and sulphates.

5      Suitable prodrugs are well-known in the art and include N-acyl derivatives, for example at any of the nitrogens in the compounds of formula (I), for example simple acyl derivatives such as acetyl, propionyl and the like or groups such as R-O-CH<sub>2</sub>-nitrogen or R-O-C(O)-nitrogen.

10     As used herein, the term halogen atom includes fluorine, chlorine, bromine or iodine.

15     The term C<sub>1-4</sub>alkyl as used herein includes straight chained and branched alkyl groups containing 1 to 4 carbon atoms, and in particular includes methyl and isopropyl.

20     The term saturated 5- or 6 - membered heterocycle containing one or two nitrogen heteroatoms as used herein includes 5- or 6- membered heterocycles containing at least one nitrogen heteroatom, and preferably two nitrogen heteroatoms, which heterocycle can be further substituted with one or more C<sub>1-4</sub> alkyl groups. A particularly suitable heterocycle is a pyrrolidine or a piperazine ring.

25     R<sup>1</sup> is aptly selected from phenyl substituted by one or more halogen atoms. Particularly, R<sup>1</sup> represents phenyl substituted by more than 1 halogen atom, such as di- or tri-halogenated phenyl. Preferably, the halogen atoms are all identical. Preferably, the halogen substituents in R<sup>1</sup> are chloro. Suitably R<sup>1</sup> is selected from 2,3,5-trichlorophenyl, 2,3-dichlorophenyl, 2,5-dichlorophenyl. Preferably, R<sup>1</sup> is 2,3,5-trichlorophenyl.

30

R<sup>3</sup> is preferably -NH<sub>2</sub>.

When R<sup>4</sup> is the group -CXNR<sup>a</sup>R<sup>b</sup> or -CXNH-(CH<sub>2</sub>)<sub>y</sub>-NR<sup>a</sup>R<sup>b</sup> where R<sup>a</sup> and R<sup>b</sup>, together with the nitrogen atom to which they are attached, form a saturated 5- or 6 - membered heterocycle containing one or two nitrogen heteroatoms atoms, this saturated 5- or 6-membered heterocycle is suitably a pyrrolidine or piperazine ring.

X is preferably =O.

The integer y is preferably 2.

When R<sup>4</sup> is the group -CXNH-(CH<sub>2</sub>)<sub>y</sub>-NR<sup>a</sup>R<sup>b</sup>, R<sup>a</sup> and R<sup>b</sup> are preferably C<sub>1-4</sub> alkyl. Preferred values for -CXNH-(CH<sub>2</sub>)<sub>y</sub>-NR<sup>a</sup>R<sup>b</sup> include for example -CONH-(CH<sub>2</sub>)<sub>2</sub>-N(CH<sub>3</sub>)<sub>2</sub>.

R<sup>4</sup> is preferably the group -CXNR<sup>a</sup>R<sup>b</sup>. Preferably X is =O and, further preferred, are compounds where R<sup>a</sup> and R<sup>b</sup> are selected from hydrogen and C<sub>1-4</sub> alkyl. Preferably R<sup>4</sup> is -CONH<sub>2</sub>, -CONH(CH<sub>3</sub>), -CONH(CH<sub>2</sub>CH<sub>3</sub>), -CONH[CH(CH<sub>3</sub>)<sub>2</sub>] or -CON(CH<sub>3</sub>)<sub>2</sub>. A particularly preferred R<sup>4</sup> is -CONH<sub>2</sub>.

It is to be understood that the present invention encompasses all isomers of the compounds of formula (I) and their pharmaceutically acceptable derivatives, including all geometric, tautomeric and optical forms, and mixtures thereof (e.g. racemic mixtures).

Preferred compounds of the present invention include:

- 5-Carboxamido-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine
- 2,6-Diamino-5-N-methylcarboxamido-3-(2,3,5-trichlorophenyl)pyrazine
- 2,6-Diamino-5-N-ethylcarboxamido-3-(2,3,5-trichlorophenyl)pyrazine
- 2,6-Diamino-5-N-isopropylcarboxamido-3-(2,3,5-trichlorophenyl)pyrazine
- 2,6-Diamino-5-N, N-dimethylcarboxamido-3-(2,3,5-trichlorophenyl)pyrazine

2,6-Diamino-5-thiocarboxamido-3-(2,3,5-trichlorophenyl)pyrazine  
and pharmaceutically acceptable derivatives thereof.

A particularly preferred compound according to the invention is:

5 5-Carboxamido-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine

and pharmaceutically acceptable derivatives thereof.

10 It is to be understood that the present invention covers all combinations of  
particular and preferred groups as described herein above.

The compounds of formula (I) are particularly useful as anticonvulsants. They  
are therefore useful in treating epilepsy. They may be used to improve the  
condition of a host, typically a human being, suffering from epilepsy. They may  
15 be employed to alleviate the symptoms of epilepsy in a host. "Epilepsy" is  
intended to include the following seizures:- simple partial seizures, complex  
partial seizures, secondary generalised seizures, generalised seizures including  
absence seizures, myoclonic seizures, clonic seizures, tonic seizures, tonic  
clonic seizures and atonic seizures.

20

The compounds of formula (I) are additionally useful in the treatment of bipolar  
disorder, alternatively known as manic depression. Type I or II bipolar disorder  
may be treated. The compounds of formula (I) may thus be used to improve the  
condition of a human patient suffering from bipolar disorder. They may be used  
25 to alleviate the symptoms of bipolar disorder in a host. The compounds of  
formula (I) may also be used in the treatment of unipolar depression.

The compounds of formula (I) are also useful as analgesics. They are therefore  
useful in treating or preventing pain. They may be used to improve the condition  
30 of a host, typically a human being, suffering from pain. They may be employed

to alleviate pain in a host. Thus, the compounds of formula (I) may be used as a pre-emptive analgesic to treat acute pain such as musculoskeletal pain, post operative pain and surgical pain, chronic pain such as chronic inflammatory pain (e.g. rheumatoid arthritis and osteoarthritis), neuropathic pain (e.g. post herpetic neuralgia, trigeminal neuralgia and sympathetically maintained pain) and pain associated with cancer and fibromyalgia. The compounds of formula (I) may also be used in the treatment or prevention of pain associated with migraine.

The compounds of formula (I) are further useful in the treatment of functional bowel disorders which include non-ulcer dyspepsia, non-cardiac chest pain and in particular irritable bowel syndrome. Irritable bowel syndrome is a gastrointestinal disorder characterised by the presence of abdominal pain and altered bowel habits without any evidence of organic disease. The compounds of formula (I) may thus be used to alleviate pain associated with irritable bowel syndrome. The condition of a human patient suffering from irritable bowel syndrome may thus be improved.

The compounds of formula (I) may also be useful in the treatment of neurodegenerative diseases, such as Alzheimer's disease, ALS, motor neuron disease, Parkinson's disease, macular degeneration and glaucoma. The compounds of formula (I) may also be useful in neuroprotection and in the treatment of neurodegeneration following stroke, cardiac arrest, pulmonary bypass, traumatic brain injury, spinal cord injury or the like.

The compounds of formula (I) are further useful in the treatment of tinnitus.

Still further, the compounds of formula (I) are also useful in preventing or reducing dependence on, or preventing or reducing tolerance or reverse tolerance to, a dependence - inducing agent. Examples of dependence inducing



agents include opioids (eg morphine), CNS depressants (eg ethanol), psychostimulants (eg cocaine) and nicotine.

5 According to a further aspect of the invention, we provide a compound of formula (I) or a pharmaceutically acceptable derivative thereof for use in human or veterinary medicine.

10 According to another aspect of the invention, we provide use of a compound of formula (I) or a pharmaceutically acceptable derivative thereof in the manufacture of a medicament for use in the treatment of a disorder substantially as hereinbefore described.

15 According to a further aspect of the invention, we provide a method of treating a human or animal subject suffering from, or susceptible to, a disorder substantially as hereinbefore described, which method comprises administering to said subject a therapeutically effective amount of a compound of formula (I) or a pharmaceutically acceptable derivative thereof.

20 It is to be understood that reference to treatment as used herein includes treatment of established symptoms and prophylactic treatment, unless explicitly stated otherwise.

25 The compound of formula (I) and its salts may be administered orally at a dose of from 0.1 to 10 mg/kg body weight per day and more particularly 0.3 to 3 mg/kg body weight per day, calculated as the free base. The dose range for adult human beings is generally from 8 to 1000 mg/day, such as from 35 to 800 mg/day, preferably 20 to 200 mg/day, calculated as the free base.

30 The precise amount of the compounds of formula (I) administered to a host, particularly a human patient, will be the responsibility of the attendant physician.

However, the dose employed will depend upon a number of factors including the age and sex of the patient, the precise condition being treated and its severity, and the route of administration.

5 The compounds of formula (I) and their pharmaceutically acceptable derivatives are conveniently administered in the form of pharmaceutical compositions. Thus, in another aspect of the invention, we provide a pharmaceutical composition comprising a compound of formula (I) or a pharmaceutically acceptable derivative thereof adapted for use in human or veterinary medicine. Such compositions may conveniently be presented for use in conventional  
10 manner in admixture with one or more physiologically acceptable carriers or excipients.

While it is possible for the compounds of formula (I) or a pharmaceutically acceptable derivative thereof to be administered as the raw chemical, it is  
15 preferable to present it as a pharmaceutical formulation. The formulations of the present invention comprise the compounds of formula (I) or a pharmaceutically acceptable derivative thereof together with one or more acceptable carriers or diluents therefor and optionally other therapeutic ingredients. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients  
20 of the formulation and not deleterious to the recipient thereof.

The formulations include those suitable for oral, parenteral (including subcutaneous e.g. by injection or by depot tablet, intradermal, intrathecal, intramuscular e.g. by depot and intravenous), rectal and topical (including  
25 dermal, buccal and sublingual) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing into association the compound of formula (I) or a  
30 pharmaceutically acceptable acid addition salt thereof ("active ingredient") with

the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

5

Formulations of the present invention suitable for oral administration may be presented as discrete units such as capsules, cachets or tablets (e.g. chewable tablets in particular for paediatric administration) each containing a predetermined amount of the active ingredient; as a powder or granules; as a  
10 solution or a suspension in an aqueous liquid or a non-aqueous liquid; or as an oil-in-water liquid emulsion or a water-in-oil liquid emulsion. The active ingredient may also be presented as a bolus, electuary or paste.

A tablet may be made by compression or moulding, optionally with one or more  
15 accessory ingredients. Compressed tablets may be prepared by compressing in a suitable machine the active ingredient in a free-flowing form such as a powder or granules, optionally mixed with a binder, lubricant, inert diluent, lubricating, surface active or dispersing agent. Moulded tablets may be made by moulding in a suitable machine a mixture of the powdered compound moistened with an  
20 inert liquid diluent. The tablets may optionally be coated or scored and may be formulated so as to provide slow or controlled release of the active ingredient therein.

Formulations for parenteral administration include aqueous and non-aqueous  
25 sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules and vials,  
30 and may be stored in a freeze-dried (lyophilised) condition requiring only the

addition of a sterile liquid carrier, for example, water-for-injection, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

5

Formulations for rectal administration may be presented as a suppository with the usual carriers such as cocoa butter, hard fat or polyethylene glycol.

10

Formulations for topical administration in the mouth, for example buccally or sublingually, include lozenges comprising the active ingredient in a flavoured basis such as sucrose and acacia or tragacanth, and pastilles comprising the active ingredient in a basis such as gelatin and glycerin or sucrose and acacia.

15

The compounds of the invention may also be formulated as depot preparations. Such long acting formulations may be administered by implantation (for example subcutaneously or intramuscularly) or by intramuscular injection. Thus, for example, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

20

In addition to the ingredients particularly mentioned above, the formulations may include other agents conventional in the art having regard to the type of formulation in question, for example those suitable for oral administration may include flavouring agents.

25

30

The compounds of formula (I) may be used in combination with other therapeutic agents, for example other anticonvulsants. When compounds of formula (I) or pharmaceutically acceptable derivatives thereof are used in combination with other therapeutic agents, the compounds may be administered either

sequentially or simultaneously by any convenient route. The invention thus provides, in a further aspect, a combination comprising a compound of formula (I) or a pharmaceutically acceptable derivative thereof with a further therapeutic agent.

5

The combinations referred to above may conveniently be presented for use in the form of a pharmaceutical formulation and thus pharmaceutical formulations comprising a combination as defined above together with a pharmaceutically acceptable carrier or excipient comprise a further aspect of the invention. The individual components of such combinations may be administered either sequentially or simultaneously in separate or combined pharmaceutical formulations.

When a compound of formula (I) or a pharmaceutically acceptable derivative thereof is used in combination with a second therapeutic agent active against the same disease, the dose of each compound may differ from that when the compound is used alone. Appropriate doses will be readily appreciated by those skilled in the art.

Preferred unit dosage formulations are those containing an effective daily dose, as herein above recited, or an appropriate fraction thereof, of the active ingredient. Conveniently that may be from 5 mg to 1000 mg, such as from 8 mg to 1000 mg, more conveniently 35 mg to 800 mg, and most conveniently 20 to 200 mg, calculated as the free base.

25

The present invention provides a process for preparing compounds of formula (I) and pharmaceutically acceptable derivatives thereof.


Compounds of formula (I) and pharmaceutically acceptable derivatives thereof may be prepared by any method known in the art for the preparation of compounds of analogous structure.

- 5 Suitable methods for the preparation of compounds of formula (I) and pharmaceutically acceptable derivatives thereof are described below and which form a further aspect of the invention. In the formulae that follow,  $R^1$  to  $R^4$  are as defined in formula (I) above unless otherwise stated.

10 According to a first process (A), compounds of formula (I) may be prepared by interconversion, utilising other compounds of formula (I) as precursors.


Thus, compounds of formula (I) wherein X is S may be prepared from the corresponding compound of formula (I) wherein X is O, by treatment with a  
15 thiation agent, preferably Lawessons reagent. Conveniently, the reaction is effected in the presence of a solvent or solvents, such as a halogenated hydrocarbon (e.g. dichloromethane) and/or toluene and at elevated temperature, for example 100°C.

20 According to another process (B), compounds of formula (I), where  $R^4$  is -CONH<sub>2</sub>, may be prepared from a compound of formula (II)



25 by hydrolysis under suitable reaction conditions and according to conventional procedures, eg. using sulphuric acid.

According to another process (C), compounds of formula (I) where X is O may be prepared under suitable reaction conditions by reacting a compound of formula (III)




or a protected derivative thereof, where Hal (D) represents a halogen atom, suitably bromine, with a palladium catalyst, preferably palladium (II) acetate, a ferrocene, preferably bis(diphenylphosphino)ferrocene and an amine, in the presence of carbon monoxide. The reaction is carried out in a solvent, such as dimethylformamide and at elevated temperature, for example between 65°C and 125°C.

15 Compounds of formula (II) may be prepared by reacting compounds of formula (III) or a protected derivative thereof, where Hal (D) represents a halogen atom, suitably bromine, with a cyanating agent preferably with a mixture of sodium cyanide and copper (I) cyanide. The reaction is carried out in a solvent, such as dimethylformamide and at elevated temperature, for example 130°C.

20

Compounds of formula (III) may be prepared by reacting compounds of formula (IV)

14




(IV)

or a protected derivative thereof with a suitable halogenating agent, for example N-bromosuccinimide. The reaction is conveniently carried out in a suitable solvent, such as dimethylsulfoxide and below room temperature, for example 15°C.

Compounds of formula (IV) where R<sup>2</sup> represents NH<sub>2</sub> may be prepared by cyclisation and oxidation of a compound of formula (V)

10



(V)


or a salt thereof according to conventional procedures, for example by neutralising a salt of a compound of formula (V), e.g. with lithium hydroxide in a suitable solvent such as an alcohol, e.g. methanol, under which conditions spontaneous oxidation to a compound of formula (IV) occurs.

Compounds of formula (V) may be prepared by reacting compounds of formula (VI) R<sup>1</sup>C(O)H with compounds of formula (VII)

20

R<sup>3</sup>






5 or a salt thereof, in the presence of a cyanide source, for example potassium cyanide. Compounds of formula (VI), where R<sup>1</sup> is trihalo-substituted phenyl, for example 2,3,5-trichlorobenzaldehyde, are known and may be prepared according to the methods described in WO95/07877. Compounds where R<sup>1</sup> represents alternative values are either known or may be prepared according to methods known for the preparation of known compounds.

10

Compounds of formula (VII), for example aminoacetamidine, may be prepared according to known procedures, for example, those described in Chem. Berichte, 89, 1185 (1956).

15

Compounds of formula (IV) may also be prepared from compounds of formula (VIII)




20

or a protected derivative thereof where Hal (B) represents a halogen atom, suitably chloride. For example, Hal(B) may be converted to -NR<sup>b</sup>R<sup>c</sup> by reaction with an appropriate amine in a solvent, such as ethanol and at elevated temperature, for example 180°C.

25

16

A compound of formula (VIII) may suitably be prepared from a compound of formula (IX)



5


or a protected derivative thereof by reaction with a compound of formula (X)  $R^1B(OH)_2$  in the presence of a palladium catalyst, preferably tetrakis(triphenylphosphine)palladium(0). Examples of compounds of formula (X)  $R^1B(OH)_2$  include 2,3,5-trichlorobenzeneboronic acid, 2,3-dichlorobenzeneboronic acid and 2,5-dichlorobenzeneboronic acid. Appropriately, Hal(A) in above formula (IX) is more reactive than Hal(B), and suitably Hal(A) is selected from bromide and iodide, whereas Hal(B) is aptly chloride. Compounds of formula (X) are either commercially available or can suitably be prepared from commercially available benzene analogues e.g. 1-bromo-2,3-dichlorobenzene or 2-bromo-4,6-dichloroaniline as described hereinafter in greater detail in the accompanying Examples.

10

15


20

A compound of formula (IX) can be suitably prepared by further halogenating a compound of formula (XI)



or a protected derivative thereof for example by reaction with a halogenating agent, such as N-bromosuccinimide, with stirring at below room temperature, for example between -5°C and 0°C, for several hours.

- 5 A compound of formula (XI) can be prepared from a di-halo compound of formula (XII)



10

by reaction with  $R^2H$ , where Hal(B) and Hal(C) may be the same or different halogen substituents. Aptly both Hal(B) and Hal(C) are chloride. Compounds of formula (XII) are commercially available. The reaction is carried out at elevated temperature, for example 150°C.

- 15 Certain intermediates described above are novel compounds, and it is to be understood that all novel intermediates herein form further aspects of the present invention.

Conveniently, compounds of the invention are isolated following work-up in the form of the free base. Pharmaceutically acceptable acid addition salts of the  
20 compounds of the invention may be prepared using conventional means.

Solvates (e.g. hydrates) of a compound of the invention may be formed during the work-up procedure of one of the aforementioned process steps.

- 25 The following Examples which should not be construed as constituting a limitation thereto are provided to illustrate the invention.

**Intermediate 1****2,3,5-Trichlorobromobenzene**

Sodium nitrite (3.88g, 0.056mole) was added in portions to concentrated sulphuric acid (28.16ml) stirred below 10°C. A solution of 2-bromo-4,6-dichloroaniline (12g, 0.05mole, Lancaster) in glacial acetic acid (126ml) was added maintaining the temperature below 10°C. The mixture was stirred below 10°C for 1 hr and then slowly added to a stirred solution of cuprous chloride (10.11g, 0.10mole) in concentrated hydrochloric acid (101.05ml) at room temperature. The mixture was then stirred at room temperature for 17 hrs. The product was filtered, washed with water (3x50ml), dissolved in chloroform (150ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated in vacuo to give the desired product. Yield 10g (77%), M.p. 55-57°C.

**Intermediate 2****2,3,5-Trichlorobenzeneboronic acid**

A solution of 2,3,5-trichlorobromobenzene (8.60g, 0.033mole) in dry ether (33ml) and bromoethane (4.73ml, 7.31g, 0.067mole) was added dropwise to a suspension of magnesium turnings (2.80g, 0.12mole) in dry ether (21.50ml) at room temperature. The mixture was refluxed for 0.50 hr and cooled to room temperature. The mixture was then added dropwise under nitrogen to a solution of trimethylborate (5.16ml, 5.16g, 0.05mole) in dry ether (8.60ml) maintaining the temperature below -60°C. This was warmed to room temperature overnight, then cooled in an ice-bath and treated with 2M hydrochloric acid (10ml). The ether layer was separated, washed with water (2x20ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated in vacuo. The residue was triturated with 40-60°C petroleum ether, filtered and dried in vacuo. Yield 4.57g (61%), M.p. 257-260°C.

**Intermediate 3**2-Chloro-6-amino-pyrazine

5 A suspension of 2,6-dichloropyrazine (100g, 0.67mole, Lancaster) in 0.880 ammonia (500ml) was stirred and heated at 150°C in a glass lined autoclave at 20atm for 16hrs. The cooled mixture was filtered, washed well with water (200ml) and dried. The product was recrystallised from chloroform. Yield 41.98g (48%), M.p. 150-152°C.

**Intermediate 4**2-Chloro-3-bromo-6-aminopyrazine and 2-amino-3-bromo-6-chloropyrazine

15 A solution of 2-chloro-6-aminopyrazine (20g, 0.15mole) in chloroform (1940ml) was stirred at -5°C to 0°C. N-Bromosuccinimide (27.58g, 0.15mole) was added in portions maintaining the temperature between -5 and 0°C. The mixture was warmed to room temperature and stirred for 3.50 hrs. The mixture was then washed with aqueous saturated sodium bicarbonate (1x300ml), then water (1x500ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated down in vacuo. The residue was purified by 'flash chromatography' using chloroform as the eluent. Yield of 2-chloro-3-bromo-6-aminopyrazine 13.89g (43%), M.p. 146-147°C. Yield of 2-amino-3-bromo-6-chloropyrazine 4.90g (15%), M.p. 124-125°C.

**Intermediate 5****2-Amino-6-chloro-3-(2,3,5-trichlorophenyl)pyrazine**

5 A solution of 2,3,5-trichlorobenzeneboronic acid (1.62g,  $7.18 \times 10^{-3}$  mole) in absolute ethanol (2.05ml) was added slowly to a mixture of 2-amino-3-bromo-6-chloropyrazine (1g,  $5.1 \times 10^{-3}$  mole) and tetrakis(triphenylphosphine)palladium(0) (0.334g,  $2.89 \times 10^{-4}$  mole) in benzene (10.20ml)/2M aqueous sodium carbonate (5.50ml). The mixture was refluxed for 17 hrs. The cooled reaction mixture was evaporated in vacuo and then extracted with chloroform (50ml). The chloroform  
10 layer was washed with water (2x30ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated down in vacuo. The residue was triturated with 40-60°C petroleum ether, filtered and dried in vacuo. Yield 0.205g (14%), M.p. 211-214°C.

**Intermediate 6****2-([Cyano-(2,3,5-trichlorophenyl)-methyl]-amino)-acetamidine hydrobromide**

Aminoacetamidine dihydrobromide (162.1g, 0.774 mole) was added in portions to a solution of 2,3,5-trichlorobenzaldehyde (200.0g, 0.851 mole) in methanol  
20 (2.43 litres) at room temperature. Once the addition was complete potassium cyanide (50.4g, 0.774 mole) was added in one portion to the resulting mixture. The suspension was then stirred at 25°C for 4 hours before being warmed to 50°C. The mixture is stirred at 50°C for 24 hours. Methanol was then removed *in vacuo*, the resulting solid was slurried in water (1.5 litres) and ethyl acetate  
25 (2.5 litres) and collected by filtration. The solid was then dried *in vacuo* at 50°C overnight to give the desired product. Yield 96.31g (33.4%),  $^1\text{H}$  nmr (d-6 DMSO)  $\delta$ /ppm 8.72 (3H, br, NH); 7.99 (1H, d,  $J$  2.3Hz, ArH); 7.79 (1H, d,  $J$  2.3Hz, ArH); 5.39 (1H, d,  $J$  10.6Hz, ArCH(CN)NH); 4.35 (1H, m, ArCH(CN)NH); 3.56 (2H, d,  $J$  6.4Hz, ArCH(CN)NHCH<sub>2</sub>C(=NH)NH<sub>2</sub>).

**Intermediate 7****2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine**

2-[[Cyano-(2,3,5-trichlorophenyl)-methyl]-amino]-acetamidine hydrobromide (95.36g, 0.256mole) was added in portions to a solution of lithium hydroxide monohydrate (16.11g, 0.384mole) in methanol (1.9litres) at room temperature. The resulting solution was stirred at room temperature for 3 hours before being evaporated to dryness *in vacuo*. The resulting solid was slurried in water (1.15litres) and collected by filtration. After drying at 50°C *in vacuo* the crude material was purified by recrystallisation from toluene to give the desired product. Yield 69.51g (93.8%), M.p. 178-180°C.

A suspension of 2-amino-6-chloro-3-(2,3,5-trichlorophenyl)pyrazine (0.3g,  $9.71 \times 10^{-4}$  mole) in absolute ethanol (4ml) and 0.880 aqueous ammonia (8.24ml) was stirred and heated in an autoclave at 180°C for 44hrs. The cooled mixture was evaporated *in vacuo*, and the residue extracted with chloroform (3x30ml). The combined chloroform extracts were dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated down *in vacuo*. The residue was purified by 'flash chromatography' using chloroform to 98:2 chloroform:methanol as the eluent. The product was triturated with 40-60°C petroleum ether, filtered and dried *in vacuo*. Yield 0.155g (56%), M.p. 178-180°C.

#### Intermediate 8

##### 5-Bromo-2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine

N-Bromosuccinimide (0.194g,  $1.09 \times 10^{-3}$  mole) was added over 20 min to a mixture of 2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.3g,  $1.04 \times 10^{-3}$  mole) in dimethylsulfoxide (10ml) and water (0.25ml) below 15°C. The resulting reaction mixture was stirred at 15°C for 1 hr, poured onto ice water (150ml) and extracted with ethyl acetate (2x75ml). The extract was then washed with 2 M sodium carbonate solution (50ml) and water (100ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated *in vacuo*. The residue was purified

by 'flash chromatography' using 5-13% ethyl acetate in cyclohexane as the eluent. Yield 0.183g (48%), M.p. 222-224°C.

### Intermediate 9

#### 5 5-Cyano-2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine

A mixture of 97% sodium cyanide (0.064g,  $1.306 \times 10^{-3}$  mole) and 90% copper(I) cyanide (0.135g,  $1.306 \times 10^{-3}$  mole) in dry dimethylformamide (5ml) was stirred and heated to 130°C. To the resulting clear solution was added 5-Bromo-2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.35g,  $0.95 \times 10^{-3}$  mole) in small portions, and the solution was maintained at 140-150°C for 16 hrs. The reaction mixture was cooled and evaporated in vacuo. The residue was extracted with ethyl acetate (100ml), washed with water (100ml) and brine (100ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated in vacuo. The residue was purified by 'flash chromatography' using 5-17% ethyl acetate in cyclohexane as the eluent. Yield 0.152g (51%), M.p. 277-279°C.

### Example 1

#### 20 5-Carboxamido-2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine

5-Cyano-2,6-Diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.1g,  $3.18 \times 10^{-4}$  mole) was dissolved in 75% sulphuric acid (8ml) and heated to 90°C for 20 min. The reaction was then cooled to 0°C and basified (dropwise) with 2 M sodium hydroxide solution. The yellow suspension was extracted with ethyl acetate (100ml), washed with brine (50ml), dried over anhydrous magnesium sulphate, filtered and the filtrate evaporated in vacuo. The residue was purified by 'flash chromatography' using 9-27% ethyl acetate in cyclohexane as the eluent. Yield 0.079g (75%), M.p. 245-247°C. *Anal.* Calcd for  $C_{11}H_8N_5Cl_3O$ : C, 39.73; H, 2.42; N, 21.06. Found: C, 39.88; H, 2.13; N, 20.53.

30

### Example 2



2,6-Diamino-5-N-Methylcarboxamido-3-(2,3,5-Trichlorophenyl)pyrazine

5-Bromo-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.5g,  $1.36 \times 10^{-3}$  mol), palladium(II)acetate ( $1.0 \times 10^{-2}$ g,  $4.45 \times 10^{-5}$  mol, 3mol%), and  
5 bis(diphenylphosphino)ferrocene ( $2.3 \times 10^{-2}$ g,  $4.15 \times 10^{-5}$  mol, 3mol%) were all added to dimethylformamide (5ml). Carbon monoxide gas and methylamine gas were continually bubbled through the solution whilst heating at 70°C, for 6 hours. The mixture was cooled, poured into water (50ml), extracted with ethyl acetate (3x30ml). The organics were combined and washed with 10% aq. citric acid  
10 solution (2x20ml), brine (20ml), dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated in vacuo. Purification was by 'flash chromatography' using 50% cyclohexane/ethyl acetate as the eluent. Yield 0.078g (16.6%), *Anal.* Calcd. for  $C_{12}H_{10}N_5Cl_3O$ : C, 41.56; H, 2.89; N, 20.20. Found: C, 41.42; H, 2.64; N, 19.87. N.m.r.( $CDCl_3$ ) $\delta$ ppm: 2.91+2.92(3H, 2xs), 7.34(1H, d), 7.49(1H, s broad), 7.57(1H, d).  
15

**Example 3**2,6-Diamino-5-N-Ethylcarboxamido-3-(2,3,5-Trichlorophenyl)pyrazine

20 5-Bromo-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.5g,  $1.36 \times 10^{-3}$  mol), palladium(II)acetate ( $3.0 \times 10^{-2}$ g,  $1.36 \times 10^{-4}$  mol, 10mol%), bis(diphenylphosphino)ferrocene ( $7.5 \times 10^{-2}$ g,  $1.36 \times 10^{-4}$  mol, 10mol%), and 2M ethylamine in tetrahydrofuran (9ml,  $1.8 \times 10^{-2}$ , 13eq) were all added to dimethylformamide (50ml). Carbon monoxide gas was bubbled through the  
25 mixture for 10 minutes which was then heated at 100°C, for 4 hours. The mixture was cooled, volatiles removed in vacuo, and the brown oil partitioned between water and diethyl ether. Aqueous was re-extracted with diethyl ether (x2). The organics were combined, filtered through a pad of J2 harborlite, washed with 10% aq. citric acid solution, brine, dried over anhydrous magnesium sulfate,  
30 filtered and the filtrate evaporated in vacuo. Purification was by 'flash chromatography' using 1:1 cyclohexane/ethyl acetate as the eluent. Yield 0.048g

(9.8%), N.m.r.(CDCl<sub>3</sub>) $\delta$ ppm: 1.22(3H, t), 3.41(2H, m), 4.60(2H, broad s), 7.35(1H, d), 7.42(1H, s broad), 7.57(1H, d). LC/MS EI M<sup>+</sup>= 360/362.

#### Example 4

5     2,6-Diamino-5-N-Isopropylcarboxamido-3-(2,3,5-Trichlorophenyl)pyrazine

5-Bromo-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.5g, 1.36x10<sup>-3</sup>mol), palladium(II)acetate (3.0x10<sup>-2</sup>g, 1.36x10<sup>-4</sup>mol, 10mol%), bis(diphenylphosphino)ferrocene (7.5x10<sup>-2</sup>g, 1.36x10<sup>-4</sup>mol, 10mol%), and isopropylamine (5ml, 5.9x10<sup>-2</sup>, 43eq) were all added to dimethylformamide (50ml) in an autoclave. Carbon monoxide gas was bubbled through the mixture for 10 minutes, the autoclave was then sealed and heated at 120°C, for 4 hours. The mixture was cooled, more palladium(II)acetate (3.0x10<sup>-2</sup>g, 1.36x10<sup>-4</sup>mol, 10mol%), bis(diphenylphosphino)ferrocene (7.5x10<sup>-2</sup>g, 1.36x10<sup>-4</sup>mol, 10mol%) and isopropylamine (4ml, 4.7x10<sup>-2</sup>, 34.5eq) were added and more carbon monoxide gas was bubbled through the mixture for a further 10 minutes. The autoclave was then sealed and heated at 120°C, for a further 4 hours. The mixture was then cooled, volatiles removed in vacuo, and the brown oil partitioned between water and diethyl ether. Aqueous was re-extracted with diethyl ether (x2). The organics were combined, filtered through a pad of J2 harborlite, washed with 10% aq. citric acid solution, brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated in vacuo. Purification was by 'flash chromatography' using 1:1 cyclohexane/ethyl acetate as the eluent. Yield 0.098g (19.3%), N.m.r.(CDCl<sub>3</sub>) $\delta$ ppm: 1.23(6H, d), 4.16(1H, m), 4.59(2H, broad s), 7.24(1H, broad s), 7.34(1H, d), 7.57(1H, d). LC/MS EI M<sup>+</sup>= 374/376.

#### Example 5

30     2,6-Diamino-5-N,N-Dimethylcarboxamido-3-(2,3,5-Trichlorophenyl)pyrazine

5-Bromo-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine (0.5g, 1.36x10<sup>-3</sup>mol), palladium(II)acetate (2.3x10<sup>-2</sup>g, 1.02x10<sup>-4</sup>mol, 7.6mol%) and

bis(diphenylphosphino)ferrocene ( $4.6 \times 10^{-2}$ g,  $8.30 \times 10^{-5}$ mol, 6.1mol%) were all added to dimethylformamide (50ml). Carbon monoxide gas was bubbled through for 10 minutes and then methylamine gas was continually bubbled through the solution whilst heating at 70°C, for 4.5 hours. The mixture was cooled, more  
5 palladium(II)acetate ( $2.3 \times 10^{-2}$ g,  $1.02 \times 10^{-4}$ mol, 7.6mol%) and bis(diphenylphosphino)ferrocene ( $4.6 \times 10^{-2}$ g,  $8.30 \times 10^{-5}$ mol, 6.1mol%) were added and more carbon monoxide gas was bubbled through the mixture for a further 10 minutes. Methylamine gas was continually bubbled through the solution whilst heating at 70°C, for a further 3 hours. The mixture was cooled,  
10 more palladium(II)acetate ( $2.3 \times 10^{-2}$ g,  $1.02 \times 10^{-4}$ mol, 7.6mol%) and bis(diphenylphosphino)ferrocene ( $4.6 \times 10^{-2}$ g,  $8.30 \times 10^{-5}$ mol, 6.1mol%) were added and more carbon monoxide gas was bubbled through the mixture for a further 10 minutes. Methylamine gas was continually bubbled through the solution whilst heating at 70°C, for a further 2.5 hours. The mixture was then  
15 cooled, volatiles removed in vacuo, and the brown oil partitioned between water and ethyl acetate. Aqueous was re-extracted with ethyl acetate(x2). The organics were combined, washed with 10% aq. citric acid solution, brine, dried over anhydrous magnesium sulfate, filtered and the filtrate evaporated in vacuo. Purification was by 'flash chromatography' using 2:1-1:1 cyclohexane/ethyl  
20 acetate as the eluent. Yield 0.061g (12.5%), N.m.r.(CDCl<sub>3</sub>) $\delta$ ppm: 3.07+3.24(2x3H, 2xs), 4.45(2H, broad s), 6.06(2H, broad s), 7.30(1H, d), 7.54(1H, d). LC/MS EI M<sup>+</sup>= 360/362.

### Example 6

#### 25 2,6-Diamino-5-Thiocarboxamido-3-(2,3,5-trichlorophenyl)pyrazine

5-Carboxamido-2,6-diamino-3-(2,3,5-trichlorophenyl)pyrazine, (0.05g,  $1.50 \times 10^{-4}$  mole) was dissolved in toluene (20ml) and dichloromethane (5ml). Lawessons reagent (0.061g,  $1.50 \times 10^{-4}$  mole) was added and the reaction stirred at 100°C  
30 under nitrogen for 16 hrs. A further three equivalents of Lawessons reagent (0.183g) was then added and the reaction was stirred for a further six hours. The

reaction mixture was cooled and evaporated in vacuo. The residue was purified by 'flash chromatography' using 5-22% ethyl acetate in cyclohexane as the eluent. Yield 0.009g (17%)

5	Mass Spec: (electrospray)	350 (MH <sup>+</sup> )
	Retention Time	3.23 minutes
	Micromass Platform Series 2	
	5 min Grad. (2mmABZ)	
	Instrument : Red	Flow rate : 0.8ml/min
10	Eluents : A – 0.1%V/V Formic Acid + 10mmol Ammonium Acetate	
	B – 95% MeCN + 0.05% V/V Formic Acid	
	Column : 5cm x 2.1mm ID ABZ+PLUS	
	Inject Vol: 5ul	
	Temp : RT	
15	Time	A% B%
	0.00	100 0
	3.50	0.0 100
	5.00	0.0 100
20	5.50	100 0

### Pharmacy Examples

25	<u>Sterile Formulations</u>	
	<u>Example A</u>	
		mg/ml
	Compound of the Invention	0.1 mg
	Sodium Chloride USP	9.0 mg
30	Water for Injection USP qs to	1 ml

The components are dissolved in a portion of the water for injections and the solution made up to a final volume to provide 0.1 mg/ml of the compound of the Invention. Where a salt of the compound is used the quantity of compound is increased to provide 0.1 mg/ml of the free base. The solution may be packaged for injection, for example by filling and sealing into ampoules, vials or syringes. These may be aseptically filled and/or terminally sterilised by, for example, autoclaving at 121°C.

Further sterile formulations may be prepared in a similar manner to obtain alternative concentrations of the compound.

#### Example B

	mg/ml
Compound of the Invention	0.5 mg
Mannitol	50.0 mg
Water for Injections qs to	1.0 ml

Dissolve the components in a portion of the Water for Injections. Make up to final volume and mix until homogeneous. Filter formulation through a sterilising filter and fill into glass vials. Lyophilise and seal vials. Reconstitute with appropriate solvent prior to use.

#### Formulations for Oral Administration

Tablets may be prepared by the normal methods such as direct compression or wet granulation. The tablets may be film coated with suitable film forming materials, such as an Opadry, using standard techniques. Alternatively the tablets may be sugar coated.

#### Example C

##### Direct Compression Tablet

	mg/Tablet
Compound of the Invention	5.0 mg
Magnesium Stearate	4.0 mg
Microcrystalline Cellulose( Avicel PH102) qs to	400.0 mg

5 The compound of the Invention is passed through a 30 mesh sieve and blended with the Avicel and Magnesium Stearate. The resultant blend is compressed into tablets using a suitable tablet press fitted with 11.0 mm diameter punches so as to provide 5 mg of the Compound of the Invention per tablet. Tablets of other strengths, containing for example 25 or 100 mg /tablet of the Compound of the Invention may be prepared in a similar manner.

#### Example D

##### 10 Wet Granulation Tablet

	mg/Tablet
Compound of the Invention	5.0 mg
Pregelised Starch	28.0 mg
Sodium Starch Glycollate	16.0 mg
15 Magnesium Stearate	4.0 mg
Lactose qs	400.0 mg

20 The Compound of the Invention, Lactose, Pregelised Starch and Sodium Starch Glycollate is dry mixed and then granulated using a suitable volume of Purified Water. The resultant granules are dried and then blended with the Magnesium Stearate. The dried granules are compressed using a suitable tablet press fitted with 11.0 mm diameter punches so as to provide 5 mg of the Compound of the Invention per tablet.

25 Tablets of other strengths such as 25 and 100 mg/tablet may be prepared.

#### Example E

##### Hard Gelatin Capsule

	mg/capsule
30 Compound of the Invention	5.0 mg
Microcrystalline Cellulose (Avicel PH102) qs	700.0 mg

The Compound of the Invention is passed through a 30 mesh sieve and then blended with the Microcrystalline Cellulose to provide an homogeneous blend.

The blend may then be filled into size 0EL hard gelatin capsule shells so as to provide capsules containing 5.0 mg/capsule of Compound of the Invention.

Alternative strengths such as 25 or 100mg/capsule of Compound of the Invention may be made in a similar manner.

5

#### Example F

##### Soft Gelatin Capsule

	mg/capsule
Compound of the Invention	10.0 mg
10 Polyethylene Glycol	90.0 mg
Propylene Glycol	qs 200.0 mg

Blend together the Polyethylene Glycol and Propylene Glycol using heat as necessary. Stir until homogeneous. Add the Compound of the Invention and mix  
15 until homogeneous. Fill into an appropriate gelatin mass to give soft gelatin capsules containing 200 mg of the formulation, to provide 10.0 mg /capsule of the Compound of the Invention.

Alternative strengths, for example, 5 and 25 mg/capsule of the Compound of  
20 the Invention may be prepared in a similar manner.

#### Example G

##### Syrup

25	Compound of the Invention	5.0 mg
	Sorbitol Solution	1500.0 mg
	Glycerol	1000.0 mg
	Sodium Benzoate	5.0 mg
	Flavour	12.5 mg
30	Purified Water qs to	5.0 ml

The Sodium Benzoate is dissolved in a portion of the purified water and the Sorbitol Solution added. The Compound of the Invention, Flavour and Glycerol are added and mixed until homogeneous. The resultant mixture is made up to  
35 volume with the purified water.

Other Formulations5     Example H

## Suppository

	mg/suppository
Compound of the Invention	10.0 mg
Witepsol W32, hard fat     qs	2000.0 mg

10

Melt the Witepsol W32 at approximately 36°C . To a portion of this add the Compound of the Invention and blend . Incorporate the remaining melted Witepsol W32 and blend until homogeneous. Fill mould with 2000 mg of the formulation to provide 10.0 mg /suppository of the Compound of the Invention.

15

Example I

## Transdermal

Compound of the Invention	5.0 mg
Silicone Fluid	90.9 mg
Colloidal Silicone Dioxide	5.0 mg

20

Mix the silicone fluid and active together and add the colloidal silicone dioxide. The material is then dosed into a subsequently heat sealed polymeric laminate comprised of the following: polyester release liner, skin contact adhesive composed of silicone or acrylic polymers, a control membrane which is a polyolefin ( foe example polyethylene or polyvinyl acetate) or polyurethane, and an impermeable backing membrane of a polyester multilaminate.

25

Biological Data

30

**Anticonvulsant activity**

A compound of formula (I) has been shown to have anti-epileptic activity in a rodent model of generalised epilepsy, the rat maximal electroshock test (MES) which is an animal model that reflects human generalised tonic-clonic seizures.

35



For example, male Han Wistar rats (150-200grms) were dosed i.p. with a suspension of the test compound in 0.25% methylcellulose 2 hr prior to test. A visual observation was carried out just prior to testing for the presence of ataxia.

5 Using auricular electrodes a current of 200mA, duration 300millisec, was applied and the presence or absence of hind limb extension noted. A compound of formula (I) exhibited an ED<sub>50</sub> of 1.4mg/kg compared to 6.1mg/kg for lamotrigine with a therapeutic index (ratio of the ataxia ED<sub>50</sub> and MES ED<sub>50</sub>) of 21.6 compared to 3.3 for lamotrigine.


10

No apparent adverse or toxic effects were observed during the above in vivo test due to the administration of the compounds of the invention.

**Claims**

5

1. A compound of formula (I)



10

wherein

$R^1$  is phenyl substituted by one or more halogen atoms;

15

$R^2$  is  $-NH_2$ ;

$R^3$  is  $-NH_2$  or hydrogen;

$R^4$  is  $-CXNR^aR^b$ ,  $-CXNH-(CH_2)_y-NR^aR^b$ ;

20

wherein

X is  $=O$  or  $=S$ ;

y is an integer zero, 1 or 2;

25

$R^a$  and  $R^b$ , which may be the same or different, are selected from hydrogen and  $C_{1-4}$  alkyl, or together with the nitrogen atom to which they are attached, form a saturated 5- or 6 - membered heterocycle containing one or two nitrogen

heteroatoms, which heterocycle can be further substituted with one or more C<sub>1-4</sub> alkyl groups;

and pharmaceutically acceptable derivatives thereof.

5

2. A compound according to claim 2 wherein R<sup>1</sup> is 2,3,5-trichlorophenyl.

3. A compound according to claim 1 or claim 2 wherein R<sup>3</sup> is -NH<sub>2</sub>.

10

4. A compound according to any of claims 1 to 3 wherein R<sup>4</sup> is -CONH<sub>2</sub>.

5. The compound according to any of claims 1 to 4 which is 2,6-diamino-5-carboxamido-3-(2,3,5-trichlorophenyl)pyrazine.

15

6. A compound according to any of claims 1 to 5 for use in human or veterinary medicine.

20

7. The use of a compound according to any one of claims 1 to 5 in the manufacture of a medicament for use in the treatment of epilepsy, bipolar disorder or manic depression, unipolar depression, pain, functional bowel disorders, neurogenerative diseases, neuroprotection, neurodegeneration, tinnitus, or prevention or reducing dependence on, or preventing or reducing tolerance or reverse tolerance to, a dependence-inducing agent.

25

8. A method of treating a human or animal subject suffering from, or susceptible to, epilepsy, bipolar disorder or manic depression, unipolar depression, pain, functional bowel disorders, neurogenerative diseases, neuroprotection, neurodegeneration, tinnitus, or prevention or reducing dependence on, or preventing or reducing tolerance or reverse tolerance to, a dependence-inducing agent, which method comprises administering to said subject a therapeutically effective amount of a compound according to any one of claims 1 to 5.

30


9. A pharmaceutical composition comprising a compound according to any one of claims 1 to 5 in admixture with one or more physiologically acceptable carriers or excipients.

5 10. A process for preparing a compound of formula (I) and pharmaceutically acceptable derivatives thereof according to any of claims 1 to 5 which comprises:

(A) interconversion of a compound of formula (I) into another compound of formula (I); or


10

(B) reacting a compound of formula (II)



wherein R<sup>4</sup> is -CONH<sub>2</sub>, with a hydrolysing agent; or


15 (C) reacting a compound of formula (III)



or a protected derivative thereof, wherein R<sup>4</sup> is C<sub>1-4</sub> alkyl, with palladium II acetate, bis(diphenylphosphino)ferrocene and an amine; or

20 (D) reacting a compound of formula (III)

35



or a protected derivative thereof, with sodium cyanide and copper (I) cyanide;

5 and optionally converting compounds of formula (I) prepared by any one of processes (A) to (D) into pharmaceutically acceptable derivatives thereof.

## INTERNATIONAL SEARCH REPORT

International Application No.

PCT/EP 99/06248

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 7 C07D241/26 A61K31/4965

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3 575 975 A (ED. J. CRAGOE) 20 April 1971 (1971-04-20) the whole document	1,6,7,9

☐ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- \*B\* document member of the same patent family

Date of the actual completion of the international search

17 December 1999

Date of mailing of the international search report

11/01/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Francois, J

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/EP 99/06248

## Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 8  
because they relate to subject matter not required to be searched by this Authority, namely:  
Remark: Although claim 8 is directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this International application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

